

AD-A262 854

REPORT 1

Form Approved
OMB No. 0704-0168

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing existing material, gathering the data needed, and completing and reviewing the collection of information, sending the material to the Office of Management and Budget, Paperwork Project, Washington, DC 20503.

Instructions: Search for existing data sources, gathering and reviewing other aspects of this collection of information, including personal data, may be submitted to the Office of Management and Budget, Paperwork Project, Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE February 1993		3. REPORT TYPE AND DATES COVERED Professional Paper	
4. TITLE AND SUBTITLE NEW RESULTS ON THE COMPATIBILITY OF THE STANDARD CONDITIONAL LIKELIHOOD APPROACH WITH PRODUCT SPACE CONDITIONAL EVENT ALGEBRA				5. FUNDING NUMBERS PR: ZW40 PE: 0601152N WU: DN300173	
6. AUTHOR(S) I. R. Goodman				8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Command, Control and Ocean Surveillance Center (NCCOSC) RDT&E Division San Diego, CA 92152-5001				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Chief of Naval Research Independent Research Programs (IR) OCNR-10P Arlington, VA 22217-5000				11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. ABSTRACT (Maximum 200 words) <p>It is an understatement that information in general arrives in conditional form. In fact, unconditional information can always be considered as the special case where the antecedent is universal. Nevertheless, it is surprising how little progress has been made in developing a full logic of conditional expressions, compatible with traditional conditional probability techniques. Recently, this problem has begun to be addressed by a number of researchers, who have commonly agreed that, as a basis, conditional events $(a b)$, where a and b are ordinary unconditional ones lying in some sample space (boolean or sigma-algebra), should be interpreted as intervals of events (or their logical equivalence) $[a \cdot b, b' \vee a \cdot b]$, relative to the usual event or set partial order \leq. Following this a number of conditional event algebras have been derived/proposed, possessing several desirable mathematical-logical properties and relatively easy to implement. (See, e.g. Goodman, Nguyen, & Walker, <i>Conditional Inference and Logic for Intelligent Systems: A Theory of Measure-Free Conditioning</i>, North-Holland, 1991.) However, a major drawback with these algebras is that inherently they cannot be boolean. (At most, they can have an algebraic structure which is a Stone algebra—see the above reference.) This leads, in particular, to certain problems in relating this work to the standard numerically-oriented approach of conditional probability, including the modeling of random variables and higher order conditional forms.</p> <p>Recently, a breakthrough was obtained in the development of conditional event algebra, by replacing the interval of events basis by a joint countable product probability space approach. Although computationally more intense than the previous approaches, this has not only provided solutions of the above-mentioned problems of modeling, but has yielded a basic</p> <p>CONTINUED ON BACK PAGE.</p> <p>Published in <i>Proceedings 30th Annual Bayesian Research Conference</i>, February 1992.</p>					
14. SUBJECT TERMS data fusion conditional events syntactics semantics linguistic information				15. NUMBER OF PAGES	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED				18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	
19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED				20. LIMITATION OF ABSTRACT SAME AS REPORT	

93-07493

UNCLASSIFIED

21a. NAME OF RESPONSIBLE INDIVIDUAL

I. R. Goodman

21b. TELEPHONE (Include Area Code)

(619) 553-4014

21c. PROJECT NAME

Code 421

Continued from section 13.

tie-in with conditional likelihoods. That is, if a_i, b_i are ordinary events, $i = 1, \dots, n$ and conjunction \cdot refers to the product space conjunction, then for any probability measure P for which the (a_i, b_i) are mutually independent pairs, then the usual joint product conditional likelihood form can be expressed as

$$\prod_{i=1}^n P(a_i | b_i) = \hat{P}((a_1 | b_1) \cdot \dots \cdot (a_n | b_n)),$$

where \hat{P} is the extension of P to the countable product space setting. Since the problem of determining the most appropriate likelihood form for non-independent conditionals is still an open issue, does the above equation provide a new reasonable approach to the problem by using the associated conditional event algebra as in the equation? This paper also discusses a number of related issues, including bayesian updating.

DTIC QUALITY INSPECTED 4

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	20